

# **Pelvic floor repair using Manchester technique without the need for hysterectomy**

## **Patient Information Leaflet**



British Society of Urogynaecology  
Registered charity No 1143157

## About this leaflet

The information provided in this leaflet should be used as a guide. There may be some variation in how each gynaecologist performs the procedure, the care procedures on the ward immediately after your operation and the advice given to you when you get home. You should ask your gynaecologist about any concerns that you may have.

You should take your time to read this leaflet. A page is provided at the end of the leaflet for you to write down any questions you may have. It is your right to know about your planned operation/procedure, why it has been recommended, what the alternatives are and what the risks and benefits are. These should be covered in this leaflet. You may also wish to ask about your gynaecologist's personal experience and results of treating your condition.

## Benefits and risks

The success and the risks of most operations carried out to treat prolapse and incontinence have been poorly studied and so it is often not possible to define them clearly. In this leaflet risks may be referred to as common, rare etc. or an approximate level of risk may be given. Further information about risk is explained in a leaflet published by the Royal College of Obstetricians and Gynaecologists "Understanding how risk is discussed in healthcare".

<https://www.rcog.org.uk/globalassets/documents/patients/patient-information-leaflets/pi-understanding-risk.pdf>

The following table is taken from that leaflet

Verbal description <sup>a</sup>	Risk	Risk description <sup>b</sup>
Very common	1 in 1 to 1 in 10	A person in family
Common	1 in 10 to 1 in 100	A person in street
Uncommon	1 in 100 to 1 in 1000	A person in village
Rare	1 in 1000 to 1 in 10000	A person in small town
Very rare	Less than 1 in 10000	A person in large town

<sup>a</sup> EU-assigned frequency  
<sup>b</sup> Unit in which one adverse event would be expected

## British Society of Urogynaecology (BSUG) database

In order to better understand the success and risks of surgery for prolapse and incontinence the British Society of Urogynaecology has established a national database. All members of the society are asked to enter all procedures that they carry out onto the database and you may be asked to consent to this for your operation. The data collected are being used to develop an overall picture of what procedures are being performed throughout the United Kingdom together with complications and outcomes. Individual surgeons can also use it to evaluate their own practice.

## **What is pelvic floor repair using the Manchester technique without the need for hysterectomy?**

It is an operation to support the body of the womb (uterus) by shortening the cervix (neck of womb) and conserving the body of the womb. This can be combined with a repair of the vaginal walls if these are prolapsing significantly.

It is often referred to as a Manchester repair.

## **What condition does a Manchester repair treat?**

Uterine or cervical prolapse (dropped womb or cervix).

## **What is prolapse?**

The pelvic floor muscles form a sling or hammock across the pelvic floor. These muscles together with surrounding connective tissue are responsible for keeping all the pelvic organs (bladder, uterus, vagina and rectum) in place and functioning correctly.

Pelvic organ prolapse occurs when the pelvic floor muscles, their attachments and/or the surrounding connective tissue become weak or damaged allowing the uterus (womb) or the walls of the vagina to drop. This normally occurs as a result of childbirth, but is most noticeable after the menopause.

The descent of the pelvic organs tends to be worse when you are tired or straining to empty bowels.

The amount of prolapse varies from person to person, and individual women may have prolapse affecting the uterus only, the front wall of the vagina only, the back wall of the vagina only or any combination of these. There may be bladder and bowel symptoms as well. Additionally, the cervix (neck of the womb) may become longer allowing it to drop with only a small amount of prolapse of the rest of the uterus and a Manchester Repair is a good method of surgical repair in this situation.

## **How is a Manchester Repair done?**

- The operation can be performed with you asleep (a general anaesthetic) or awake (a spinal anaesthetic) whereby you are awake but numb from the waist down.
- The operation is done through the vagina, using absorbable stitches which take up to 3-4 months to dissolve.
- Only the cervix is shortened leaving the uterus in place.

## **Other operations which may be performed at the same time.**

- **Vaginal repairs** - Very often this procedure is combined with repair of the front and/or back vaginal wall.

- **Contenance Surgery** - sometimes an operation to treat any bothersome urinary leakage can be performed at the same time as your Manchester repair. Some gynaecologists prefer to do this as a separate procedure at a later date. You should also refer to an information leaflet about the planned additional procedure.

## Benefits

The benefits of a Manchester repair, as for all surgical interventions for pelvic organ prolapse, are to resolve the symptoms of prolapse and maintain or restore as near normal bowel and bladder function and normal intercourse, without making you worse, and hopefully be as long lasting as possible.

## Risks

### General Risks of Surgery

- **Anaesthetic risk.** This is very small unless you have specific medical conditions, such as a problem with your heart, or breathing. Smoking and being overweight also increase any risks. Manchester repair can be performed with you asleep (a general anaesthetic) or awake (a spinal anaesthetic) whereby you are awake but numb from the waist down. This will be discussed with you.
  - **What can I do?** Make the anaesthetist aware of medical conditions such as problems with your heart or breathing. Bring a list of your medications. Try to stop smoking before your operation. Lose weight if you are overweight and increase your activity.
- **Bleeding.** There is a risk of bleeding with any operation but for Manchester repair this is approximately 20% less than the alternative operation, a vaginal hysterectomy. It is rare that we have to transfuse patients after their operation.
  - **What can I do?** Please let your doctor know if you are taking a blood-thinning tablet such as warfarin, aspirin, clopidogrel or rivaroxaban as you may be asked to stop them before your operation.
- **Infection.** There is a small risk of infection with any operation (about 5 to 13 cases in 100 operations). If it occurs, an infection can be a wound infection, vaginal infection or a urinary infection, and is usually treated with antibiotics. A significant infection is rare. The risk of infection is reduced by routinely giving you a dose of antibiotic during your operation. Chest infection may also occur because of the anaesthetic.
  - **What can I do?** Treat any infections you are aware of before surgery. After surgery, regular deep breathing exercises can help prevent chest infections; the nurses will guide you how to do this.

- **Deep Vein Thrombosis (DVT).** This is a clot in the deep veins of the leg. Occasionally this clot can travel to the lungs (pulmonary embolism) which can be very serious and in rare circumstances it can be fatal (less than 1 in 100 of those who get a clot). The risk increases with obesity, severe varicose veins, infection, immobility and other medical problems. The risk is significantly reduced by using special stockings and injections to thin the blood.
  - **What can I do?** Stop taking any hormones such as hormone replacement therapy (HRT) and some types of birth-control pills 4 weeks before surgery. These can usually be restarted 4 weeks following surgery when the risk of blood clots has reduced. Do not arrange surgery the day after a long car journey or flight. As soon as you are awake start moving your legs around. Keep mobile once you are at home and continue to wear your compression stockings during times when you are less mobile.
- **Wound complications.** The wound within the vagina can become infected or occasionally stitches can become loose allowing the wound to open up or tighten causing discomfort.
  - **What can I do?** Keep the surrounding area clean and dry carefully after washing using a clean towel or a hairdryer on a cool setting. Do not douche the vagina or use tampons

### General risks of prolapse surgery

- **Getting another prolapse.** There is little published evidence of exactly how often prolapse recurs. Recurrence of the same prolapse probably occurs in about 1 in 10 cases but it is generally believed that about 3 in 10 women who have an operation for prolapse will eventually require treatment for another prolapse. This is because the vaginal tissue is weak. For women having a Manchester repair however, it seems that fewer women will need repeat surgery for prolapse than will be the case for those having a vaginal hysterectomy to treat a prolapse of the uterus. Sometimes even though another prolapse develops it is not bothersome enough to require further treatment.
  - **What can I do?** Keeping your weight normal for your height (normal BMI), avoiding unnecessary heavy lifting, and not straining on the toilet, may help prevent a further prolapse, although even if you are very careful it does not always prevent it.
- **Failure to cure** symptoms. Even if the operation cures your prolapse it may fail to improve your symptoms.
- **Overactive bladder symptoms** (urinary urgency and frequency with or without incontinence) usually improve after the operation, but occasionally can start or worsen after the operation.
  - **What can I do?** If you experience this, please let your doctor know so that treatment can be arranged.

- **Stress incontinence** Having a prolapse sometimes causes some kinking of the tube through which you pass urine (urethra). This can be enough to stop urine leaks on coughing, laughing or sneezing. By correcting the prolapse this kink gets straightened out and the leaks are no longer stopped. It is difficult to define an exact risk but it is reported to be in the order of 10% (1 in 10).
  - **What can I do?** Doing pelvic floor exercises regularly can help to prevent stress incontinence.
- **Bladder emptying or voiding problems** generally improve after surgery for prolapse but there may be problems emptying the bladder in the first few days. Your doctor may wish to do bladder tests (urodynamics) prior to surgery to predict post-operative voiding difficulties. There can be persistence of voiding problems in 1 in 10 women.
  - **What can I do?** If you experience difficulty passing urine, you may wish to lean forwards or even stand slightly to allow better emptying of your bladder. Make sure that you have your legs apart than having your knees together when sitting on the toilet. Waiting for two minutes after the initial void and trying again may help. This is known as the double void technique.
- **A change in the way your bowel works.** Some patients experience worsening constipation following surgery. This may resolve with time. It is important to try to avoid being constipated following surgery to reduce prolapse recurrence.
  - **What can I do?** If you are struggling with constipation after simple changes in diet and fluid intake, your doctor/GP may prescribe some laxatives.
- **Painful sexual intercourse.** Once the wound in the vagina has healed which usually takes about 6 weeks it is safe to have intercourse. Some women find sex is uncomfortable at first but it gets better with time. Occasionally pain on intercourse can be long-term or permanent.
- **Altered sensation during intercourse:** Sometimes the sensation during intercourse may be less and occasionally orgasm may be less intense. On the other hand repair of your prolapse may improve it.

### Specific risks of Manchester Repair

- As in all operations for pelvic organ prolapse the surgery is close to the bowel and the bladder, so injuries to these structures happen occasionally, but can normally be repaired straight away without long term consequence. They may increase the time you need in hospital to recover.
- After the Manchester repair, the shortened cervix can become narrowed if skin heals over it, which can stop period blood being released – this is rare.

## Before the operation - Pre-op assessment

Usually you are seen in a preoperative clinic some weeks before your planned operation. At that visit you will be seen by a nurse and possibly also a doctor. You will be asked about your general health and any medications you take. Your blood pressure will be checked and you may have tests to assess your heart and breathing. Blood tests will be taken to check you for anaemia and other things according to your medical condition. Swabs may be taken from your nose and groin to make sure that you do not carry MRSA (bacteria that are very resistant to antibiotics and may cause problems after your operation). You may be asked to sign a consent form if this has not been done already.

## After the operation - in hospital

- **Pain relief.** Pain can be controlled in a number of ways depending on the preference of your anaesthetist and/or gynaecologist. Options are an epidural, injection of local anaesthetic into the tissues during the operation, self-administration of pain relief (patient controlled analgesia - PCA), drugs in a drip, tablets or suppositories. A Manchester repair is not a particularly painful operation but sometimes you may require tablets or injections for pain relief. Some women describe the pain as similar to a period. It is often best to take the pain killers supplied to you on a regular basis aiming to take a pain killer before the pain becomes a problem
- **Drip.** This is to keep you hydrated until you are drinking normally. The drip is usually removed within 24 hours.
- **Catheter.** You may have a tube (catheter) draining the bladder. The catheter may give you the sensation as though you need to pass urine but this is not the case. It is usually removed the morning after surgery or sometimes later the same day.
- **Pack.** Some gynaecologists insert a length of gauze into the vagina at the end of the operation. It acts as pressure bandage to control bleeding and maintains the shape of the vagina and is usually removed 24 to 48 hrs after surgery.
- **Vaginal bleeding.** There may be slight vaginal bleeding like the end of a period after the operation.
- **Eating and drinking.** You can drink fluids soon after the operation and will be encouraged to start eating as soon as tolerated.
- **Preventing DVT (deep vein thrombosis).** The same day or the day after your operation, you will be encouraged to get out of bed and take short walks around the ward. This improves general wellbeing and reduces the risk of clots in the

legs. You may be given a daily injection to keep your blood thin and reduce the risk of blood clots until you go home or longer in some cases.

- **Going home.** You are usually in hospital for one or two days. If you require a sick note or certificate please ask.

### After the operation – at home

- Mobilisation is very important; using your leg muscles will reduce the risk of clots in the back of the legs (DVT).
- Bath or shower as normal.
- Do not use tampons for 6 weeks and avoid douching the vagina
- Any of the stitches under the skin will melt away by themselves. The surface knots of the stitches may appear on your underwear or pads after about 2 weeks, this is quite normal. There may be little bleeding again after about 2 weeks when the surface knots fall off, this is nothing to worry about.
- You are likely to feel tired and may need to rest in the daytime from time to time for a month or more, this will gradually improve.
- It is important to avoid stretching the repair particularly in the first weeks after surgery. Therefore, avoid constipation and heavy lifting. The deep stitches dissolve during the first 3 months and the body will gradually lay down strong scar tissue over a few months.
- Avoiding constipation
  - Drink plenty of water / juice
  - Eat fruit and green vegetables especially broccoli
  - Plenty of roughage e.g. bran / oats
- Any constant cough is to be treated promptly. Please see your GP as soon as possible.
- At 6 weeks gradually build up your level of activity.
- After 3 months, you should be able to return completely to your usual level of activity.
- You should be able to return to a light job after about 6 weeks, a busy job in 12 weeks. Avoiding all unnecessary heavy lifting will possibly reduce the risk of the prolapse recurring.
- You can drive as soon as you can operate the pedals and look over your shoulder without discomfort, generally after three weeks, but you must check this



with your insurance company, as some of them insist that you should wait for six weeks.

- You can start having sex whenever you feel comfortable enough after about 6 weeks. You will need to be gentle and may wish to use lubrication
- You usually have a follow up appointment anything between 6 weeks and 6 months after the operation. This maybe at the hospital (doctor or nurse), with your GP or by telephone. Sometimes follow up is not required.
- See link: <https://www.rcog.org.uk/globalassets/documents/patients/patient-information-leaflets/recovering-well/pelvic-floor-repair-operation.pdf>

### **What to report to your doctor after surgery**

- Heavy vaginal bleeding
- Smelly vaginal discharge
- Severe pain
- High fever
- Pain or discomfort passing urine or blood in the urine
- Difficulty opening your bowels.
- Warm, painful, swollen leg
- Chest pain or difficulty breathing

## Treatment Alternatives

### Non-surgical

- **Do nothing.** If the prolapse is not too bothersome treatment is not necessarily needed. If, however, the prolapse permanently protrudes through the opening to the vagina and is exposed to the air, it may become dried out and eventually ulcerate. Even if it is not causing symptoms in this situation it is probably best to push it back with a ring pessary (see below) or have an operation to repair it. Weight reduction in overweight women and avoiding risk factors such as smoking (leading to chronic cough), heavy weight lifting jobs and constipation may help with symptom control. The prolapse may become worse with time but it can then be treated.
- **Pelvic floor exercises (PFE).** The pelvic floor muscles support the pelvic organs. Strong muscles can help to prevent a prolapse dropping further. PFE are unlikely however to provide significant improvement for a severe prolapse protruding outside the vagina. A women's health physiotherapist can explain how to perform these exercises with the correct technique. It is important that you try the above to help to manage the symptoms of your prolapse and to prevent it becoming worse. It is also very important to continue with your pelvic floor exercises even if you have opted for other treatment options.
- **Pessary.** A vaginal device, a pessary (see image below), may be placed in the vagina to support the vaginal walls and uterus. A pessary is usually used continuously and changed by a doctor or nurse every 4 to 12 months depending upon the type used and how well it suits you. Alternatively, if you prefer, you may be taught to replace the pessary yourself. It is possible to lead a normal life with continuation of activities such as bathing, cycling, swimming and, in some cases, sexual intercourse. Ongoing care is often at the GP practice but some women will need to be kept under review in the Gynaecology clinic. Pessaries are very safe and many women to choose to use one long term rather than have an operation. On occasions their use has to be discontinued due to bleeding, discharge, sexual difficulties or change in bladder function but these all stop quickly after removal. Sometimes it will take several visits to the clinic to determine the best size for you. A pessary is not suitable for all women.



## Surgical

The following table lists the different operations that can be considered to treat uterine prolapse. Further information on the operations is available in separate leaflets. All operations are not available in all hospitals. Your consultant may recommend a particular operation depending on his or her preference and expertise, or your individual needs.

Surgical Treatment	Advantages	Disadvantages
<b>Manchester repair</b> (removal of cervix only via the vagina) (described in this leaflet)	No abdominal incision(s) Main body of uterus still present so pregnancy is possible. Can be done with you awake or asleep	Rarely stenosis (narrowing) of cervix causes pain Pregnancy can be complicated by premature labour
<b>Vaginal Hysterectomy</b> (removal of uterus via the vagina)	No abdominal incision(s) Uterus removed so no risk of cancer of cervix or uterus in future. Can be done with you awake or asleep	Risk of prolapse of the vault (top) of the vagina in the future
<b>Sacrohysteropexy</b> - laparoscopic (key hole) or abdominal (open operation)	Mesh provides strong and continuing support to the uterus reducing the chance of prolapse recurrence. May also treat a co-existing vaginal prolapse. No cuts or stitches in vagina. Vaginal length maintained. Uterus still present so pregnancy is possible. Minimal blood loss and shorter length of hospital stay (equivalent to other options) with laparoscopic approach.	Requires a general anaesthetic (asleep) for laparoscopic or open surgery As mesh is used there is a small risk that the mesh will work its way into surrounding tissues. <b>Only if open surgery</b> <ul style="list-style-type: none"> <li>• More painful than the other procedures</li> <li>• Slower return to normal activities</li> <li>• Longer hospital stay</li> </ul>
<b>Vaginal Sacrospinous Hysteropexy</b> (stitches to support womb inserted through vagina)	No abdominal incision(s) Pregnancy still possible although prolapse might recur during or after pregnancy Can be done with you awake or asleep	Can cause temporary buttock pain Variable long-term success with recurrence of uterine prolapse 14-30%.
<b>Colpocleisis</b> (closing of vagina)	High success rates (90-95%) both for prolapse of the uterus and the walls of the vagina. No abdominal incision(s) Can be done with you awake or asleep	Sexual intercourse will never be possible after this operation. Not possible to take a smear Difficult to investigate inside the uterus if abnormal bleeding occurs Urinary incontinence in the future may be more difficult to treat

## More information

If you would like to know more about uterine prolapse and the treatments available for it, you may try the following sources of information.

- Ask your GP.
- Ask the Doctor or Nurse at the hospital.
- Look at a website such as
  - NHS choices at <http://www.nhs.uk/pages/home.aspx>
  - Patient UK at <http://patient.info/health>
  - Royal College of Obstetricians and Gynaecologists Recovering Well leaflet (no specific leaflet for Manchester Repair ) at <https://www.rcog.org.uk/globalassets/documents/patients/patient-information-leaflets/recovering-well/pelvic-floor-repair-operation.pdf>
  - Royal College of Obstetricians and Gynaecologists patient information leaflet – Pelvic organ prolapse at <https://www.rcog.org.uk/globalassets/documents/patients/patient-information-leaflets/gynaecology/pi-pelvic-organ-prolapse.pdf>
  - Patient information leaflets for your own hospital and others (usually available on line)

## Acknowledgements

Mr. Clive Spence-Jones, BSUG patient information committee project lead for this leaflet, on behalf of BSUG.

Miss Farah Lone, Consultant Urogynaecologist, Royal Cornwall Hospitals for the photograph of vaginal pessaries

## Making a decision - things I need to know before I have my operation.



**Shared Decision Making**

if you are asked to make a choice, you may have lots of questions that you want to ask. You may also want to talk over your options with your family or friends. It can help to write a list of the questions you want answered and take it to your appointment.

**Ask 3 Questions**

To begin with, try to make sure you get the answers to three key questions if you are asked to make a choice about your healthcare.

1. What are my options?
2. What are the pros and cons of each option for me?
3. How do I get support to help me make a decision that is right for me?

These resources have been adapted with kind permission from the MAGIC programmes, supported by the Health Foundation

\* Ask 3 Questions is based on Shephard H., et al. Three questions that patients can ask to improve the quality of information physicians give about treatment options: A cross-over trial. Patient Education and Counselling, 2011; 84: 379-85

AQUA Advancing Quality Alliance    NHS    <http://www.advancingqualityalliance.nhs.uk/SDM/>

**Please list below any questions you may have, having read this leaflet.**

1).....

2).....

3).....

**Please describe what your expectations are from surgery.**

1).....

2).....

3).....